

Strataderm®

scar therapy gel




Strataderm is a self-drying gel for the management of **NEW and OLD scars**.

- Used to soften and flatten raised scars
- Used to relieve itching and discomfort
- Used to reduce redness and discoloration

Stratacel®

film-forming wound dressing for sensitive skin



Post-treatment



post-treatment



Day 11



Post-procedure



Day 2



Day 7

Stratamed is a self-drying **FULL CONTACT** film forming gel for optimal healing.
Gives patients a faster recovery time and the ability to wear sunscreen and make-up immediately post laser!

**Ask your doctor about advanced
wound healing and professional scar
management today!**


 **Stratapharma**
Switzerland



Strataderm®

scar therapy gel

Strataderm – for the prevention and treatment of hypertrophic scars and keloids



Scars, often viewed as aesthetically displeasing, are among the most common and frustrating problems after injury.^{1,2}

- Silicone gel is internationally recommended as first-line treatment in scar management^{3,4}
- Strataderm can be used as monotherapy, as well as in combination with more invasive scar therapy options⁴

For scars old and new

- Prevents abnormal and excessive scar formation⁵
- Softens and flattens raised scars⁶
- Relieves itching and discomfort⁶
- Reduces redness and discoloration⁶

Convenient and easy to use

- Rapidly drying, non-sticky and waterproof
- Can be used on joints and areas with contours
- Transparent, odorless and can be covered with make-up or sunscreen



Linear hypertrophic scar (after nevis excision)⁷

14 months after excision



Monotherapy treatment with Strataderm, left scar remained untreated

After 2 months treatment



Strataderm is semi-occlusive, which allows the skin to breathe and remain hydrated. This helps to achieve a softer, smoother and flatter scar.^{5,8}



Strataderm forms a durable, flexible and waterproof sheet⁷, which protects scars from microbial and chemical invasion and helps normalizing collagen production.⁸



Strataderm modulates the level of basic fibroblastic growth factors (bFGF), which prevents the formation of hypertrophic scar tissue.^{2,8}



Stratapharma
Switzerland

CE Class I Medical Device,
FDA/TGA Registered

www.strataderm.com

5g \$22.95 (only available online), 10g \$34.95, 20g \$54.95

References:

1. Atiyeh BS. Aesth Plast Surg 2007;31:468-492;
2. Hanasano MM et al. Facial Plast Surg 2004;6:88-93;
3. Mustoe TA et al. Plast Reconstr Surg 2002; 110:560-571;
4. Gold MH et al. Dermatol Surg 2014;40:817-824;
5. Monstrey S et al. JPRAS 2014;1-9;
6. Tobola J. Dermatol Estetyczna 2004;35(6):323-328;
7. Data on file. Stratapharma AG;
8. Mustoe TA. Aesth Plast Surg 2008;32:82-92