

Fox Valley Plastic Surgery, S.C. www.fvpsurgery.com
 2400 Witzel Avenue, Suite A 2500 E Capitol Drive, Suite 1500
 Oshkosh, WI 54904 Appleton, WI 54911
 920-233-1540 920-358-1810
 414-386-4625 Fax 920-358-1819 Fax

Name: _____

DOB: _____

CONSENT FOR COTININE TEST

Blood flow to tissues is always a great concern during any surgical procedure because decreased blood flow can produce dangerous complications. Blood flow to tissue may be compromised with incisions and dissection. This effect is typically controllable, but when coupled with nicotine – a vasoconstrictor which decreases blood flow – reduced blood flow may result in a healing complication. Smokers face longer healing times, blood clots, wounds that break open, and greater risk of infection.

In a breast augmentation, liposuction, or blepharoplasty, the risk of blood flow compromise is lower because the incisions are smaller. However, in a breast lift or tummy tuck, there are much longer incisions and more dissection. The decrease in blood flow to the tissues, combined with the constricting effects of nicotine, can cause parts of the breast, nipple, or abdominal tissue to die. This complication also applies to breast reconstruction, breast reduction, and facelift surgeries.

In addition to actual cigarette smokers, patients who are exposed to secondhand smoke or who use e-cigarettes fall into the smoker category. Vaporizers do not decrease the amount of nicotine in tobacco, only the amount of smoke generated. Patches also contain nicotine. Marijuana alone does not contain nicotine; however, it is sometimes combined with tobacco.

For certain surgeries, we require the cessation of smoking at least one month prior to surgery to reverse some of the effects of nicotine. You must remain off any sort of nicotine for one month following surgery to insure proper healing and an optimal outcome. You could quit a week before surgery, but it takes two weeks for cotinine (the predominant metabolite of nicotine) to clear the system.

The surgery that you have selected requires good blood flow. Smokers are required to take a nicotine test in the office prior to the actual surgery. Adequate notice of cotinine testing is always given, so there is no reason for a failed test. The blood test gives an immediate result. If you fail the nicotine test, the surgery will be cancelled, and rescheduled if you wish. This is the rescheduling/cancellation fee schedule, which also applies to failed cotinine tests.

Days Prior to Surgery	Rescheduling Fee	Cancellation Fee
Over 21 days	\$200.00	\$300.00
15-21 days	\$400.00	\$500.00
8-14 days	\$600.00	\$700.00
1-7 days	\$800.00	\$900.00
24 hours or less	\$1000.00	\$1000.00

Smoking is such a critical factor to the success of a surgery that FVPS is taking a firm position against smoking. FVPS would love to perform every scheduled surgery; we hope that no patients fail the nicotine test. Because we want the best possible surgical outcomes, we will not perform certain surgeries on patients who have nicotine in their blood.

I have read and understand this Consent for Cotinine Test. I consent to a nicotine test prior to surgery. If I test positive for nicotine, I understand that surgery will be cancelled and I will be charged a rescheduling/cancellation fee.

Patient's or Responsible Party's Signature

Date

Witness Signature

Date